

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number
WO 2004/021012 A2

(51) International Patent Classification⁷: **G01N 33/68**

(21) International Application Number:
PCT/IL2003/000706

(22) International Filing Date: 28 August 2003 (28.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/406,687 29 August 2002 (29.08.2002) US

(71) Applicant (for all designated States except US): **DIAG-
NOSTIC TECHNOLOGIES LTD.** [IL/IL]; Ha'Histadrut
Blv. 49, 31250 Haifa (IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MEIRI, Hamutal**
[IL/IL]; 4 Weiss Street Apt. 34, 69124 Tel Aviv (IL).

(74) Agent: **REINHOLD COHN AND PARTNERS**; P.O.Box
4060, 61040 Tel-Aviv (IL).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF DIAGNOSIS OF PREGNANCY-RELATED COMPLICATIONS



(57) **Abstract:** A method for diagnosing pregnancy-related complications in a pregnant woman. The method comprises (a) provid-
ing PP13-responding cells having a membrane; (b) exposing the cells to standard PP13; (c) exposing PP13-responding cells to PP13
obtained from the woman, and (d) determining the existence of a modification in the permeability of the cell membrane in (b) and
in (c) as a result of exposure to PP13. The modification in permeability in (b) and (c) is then compared, a change in the permeability
of (c) as compared to the permeability in (b) indicating the existence of a pregnancy complication in the woman.